

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/279230959>

Can Variability of Pattern ERG Signal Help to Detect Retinal Ganglion Cells Dysfunction in Glaucomatous Eyes?

Article · July 2015

DOI: 10.1155/2015/571314

CITATIONS

3

READS

76

3 authors:



Alberto Mavilio

Azienda Sanitaria Locale Brindisi

3 PUBLICATIONS 4 CITATIONS

SEE PROFILE



Francesca Scrimieri

19 PUBLICATIONS 903 CITATIONS

SEE PROFILE



Donato Errico

Ospedale Cardinale Giovanni Panico

6 PUBLICATIONS 15 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



RE-PERG a new electrophysiological procedure for glaucoma diagnosis [View project](#)

Clinical Study

Can Variability of Pattern ERG Signal Help to Detect Retinal Ganglion Cells Dysfunction in Glaucomatous Eyes?

Alberto Mavilio,¹ Francesca Scrimieri,² and Donato Errico²

¹Social Health District, Glaucoma Center, Azienda Sanitaria Locale, Via Fermi, 72015 Fasano, Brindisi, Italy

²Ophthalmology Unit, Glaucoma Service, Azienda Ospedaliera "Cardinale G. Panico", 73039 Tricase, Lecce, Italy

Correspondence should be addressed to Alberto Mavilio; a.mavilio@gmail.com

Received 29 August 2014; Revised 30 December 2014; Accepted 1 January 2015

Academic Editor: Antonio Ferreras

Copyright © 2015 Alberto Mavilio et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Objective. To evaluate variability of steady-state pattern electroretinogram (SS-PERG) signal in normal, suspected, and glaucomatous eyes. **Methods.** Twenty-one subjects with suspected glaucoma due to disc abnormalities (GS), 37 patients with early glaucoma (EG), and 24 normal control (NC) were tested with spectral-domain optical coherence tomography (SD-OCT), standard automated perimetry (SAP), and SS-PERG. Mean deviation (MD), pattern standard deviation (PSD), retinal nerve fiber layer (RNFL), and ganglionar complex cells (GCC) were evaluated. The SS-PERG was recorded five consecutive times and the amplitude and phase of second harmonic were measured. PERG amplitude and coefficient of variation of phase (CVphase) were recorded, and correlation with structural and functional parameters of disease, by means of one-way ANOVA and Pearson's correlation, was analysed. **Results.** PERG amplitude was reduced, as expression of retinal ganglion cells (RGCs) dysfunction, in EG patients and GS subjects compared to NC patients ($P < 0.0001$). CVphase was significantly increased in EG patients and GS subjects, compared to healthy ($P < 0.0001$), and it was also correlated with PSD ($P = 0.0009$), GCC ($P = 0.028$), and RNFL ($P = 0.0078$) only in EG patients. **Conclusions.** Increased intrasession variability of phase in suspected glaucomatous eyes may be a sign of RGCs dysfunction.

1. Introduction

Glaucoma is a progressive optic neuropathy characterized by death of retinal ganglion cells (RGCs), clinically manifested as typical alterations of the optic nerve head (ONH) and retinal nerve fiber layer (RNFL) correlated with visual field defects.

The standard automated perimetry (SAP) is the main tool for the detection of visual field loss. However, the subjective nature of the test and the fact that the examination reveals glaucomatous defects only when 30 to 40% of the fibers have already been lost [1, 2] have increased the interest of research towards alternative diagnostic tools.

The spectral-domain optical coherence tomography (SD-OCT), a good surrogate accepted for the diagnosis of glaucoma, has been shown to objectively measure ONH and RNFL [3–5].

Because most of the retinal ganglion cells are located in the macula, the study of this area, in particular the ganglion cells complex (GCC), has been proposed in the early

evaluation of glaucoma variations, in addition to the changes that occur in ONH and RNFL [6–10].

Pattern electroretinogram (PERG) alterations reflect the electrical activity of RGCs [11, 12] and has been widely used to detect the loss of function of RGCs in glaucoma [13, 14]. Cross-sectional studies have shown that PERG is frequently altered in glaucoma suspects (GS) and patients with early glaucoma with respect to normal controls [13–19].

PERG has been shown to be abnormal before both the occurrence of visual field defects, as measured by SAP, and RNFL loss, as assessed by OCT [13].

An optimized model of PERG for glaucoma screening (PERGLA) is a relatively new diagnostic tool and fast and user-friendly for the evaluation RGCs dysfunction [20, 21].

The steady-state PERG is recorded in response to stimulus of high temporal frequency [22]. Better than a transient stimulus (slow), a steady-state stimulus (fast) is able to show a glaucomatous dysfunction since it submits the RGCs to a greater metabolic stress [20]. The steady-state stimulus

determines a sinusoidal response that is analysed by the Fourier transform [23–25]. In this way, the second harmonic, that is, the harmonic that has a frequency twice that of the stimulus, is isolated. Two components of this harmonic, that is, amplitude and phase, show typical alterations in glaucoma. In particular, the amplitude is reduced in patients with glaucoma and ocular high pressure (OHT) compared to healthy subjects [14, 21], while the phase remains constant or at most tends to delay with age [20]. Steady-state PERG has been reported to have high test-retest repeatability [26]. However it is very important to work with a good signal-to-noise ratio (SNR) [27]. In particular the phase variability has been showed to be very limited in the retest within and between trials [16, 26].

The biological variability of a measurement is a physiological characteristic that can bias not only diagnostic imaging [28] but also the different adaptation of the bioelectrical response to an external visual stimulus [29, 30].

Variations of the phase are little affected by opacity of the media and deterioration of optics that may cause a nonspecific reduction of PERG amplitude [31].

If the variability of the phase is the expression of a dysfunction of RGCs that precedes cell death, we hypothesized that the within-trial variability of the PERG signal, individual test-retest of the same eye of early glaucoma patients, is greater than the one physiologically present in healthy individuals. Therefore, we checked if such variability correlates with markers of disease severity such as retinal thickness and visual field indices.

2. Materials and Methods

Participants were recruited from the Glaucoma Center of the Brindisi Social Health District, Mesagne, Italy. The participants were divided into 3 groups: early glaucoma (EG), glaucoma suspects (SG), and normal control (NC).

A total of eighty-two eyes were included. The criteria for classification in the EG group, in accordance with the EGS guidelines (http://www.eugs.org/eng/EGS_guidelines.asp), were as follows: appearance of the optic disc and peripapillary nerve fiber layer suspected for glaucoma damage (increased ratio cup/disc, asymmetry ratio of cup/disc, notch or narrowing of the neuroretinal rim, disc haemorrhage, and thinning of the peripapillary nerve fiber layer) or visual field suspicious for glaucomatous damage in the absence of clinical signs of other optic neuropathies (nasal step, paracentral scotoma, and altitudinal defect) or a constant elevated intraocular pressure.

The severity of glaucoma was evaluated functionally by means of SAP and anatomically by means of RNFL and GCC thickness measurement by SD-OCT.

Thirty-seven patients had early glaucoma (EG), defined as consecutive repeatable abnormal SAP results according to the normative database of the instrument; 21 had suspected glaucoma, defined as optical discs apparently abnormal (presence of thinning of the neuroretinal rim or localized or diffuse RNFL defects indicative of glaucoma as evaluated by Stereo photography of the fundus) without repeatable abnormal SAP results. 24 age-matched healthy subjects,

defined as those with IOP <22 mmHg with no history of elevated IOP, optic disc apparently healthy, and normal SAP repeatable results, were included in the control group. The EG patients were under medical topical treatment with beta-blockers, prostaglandin analogues, and alpha adrenergic receptors eye drops. Each participant of the trial underwent a comprehensive ophthalmic evaluation, including review of medical history, best-corrected visual acuity testing, IOP measuring by means of Goldmann applanation tonometry, ultrasound pachymetry (Pachmate GH55), slit lamp biomicroscopy, gonioscopy, and dilated fundus examination with a 78 D lens. All participants had best-corrected visual acuity $\geq 20/30$ (Snellen), spherical refraction within ± 5.0 D and cylinder correction within ± 2.0 D, transparent ocular media (nuclear color/opalescence, cortical or posterior subcapsular lens opacity <1) according to the Lens Opacity Classification System III, and open angle on gonioscopy. Coexisting retinal disease, diabetes, Parkinson's disease, or nonglaucomatous optic neuropathy, potentially able to determine nonspecific PERG abnormality, were excluded [32–34].

One eye per patient who met the criteria mentioned above was included in the study. When both eyes of the patient were eligible, the one with best-corrected visual acuity was selected. In case of equal visual acuity, right eye, by convention, was selected for evaluation.

This trial followed the tenets of the Declaration of Helsinki for human studies. The study was approved by Ethical Committee of the Brindisi Social Health District. Informed written consent was obtained by all subjects after the nature of the test and possible risks were explained in detail.

2.1. Spectral-Domain Optical Coherence Tomography. Peripapillary RNFL thickness was assessed by Zeiss Cirrus HD-OCT 500 (software version 7.0.1.290, Carl Zeiss Meditec, Dublin, CA). The protocol Optic Disc Cube 200 \times 200 was used to perform a circular scan 3.46 mm in diameter, which was automatically targeted around the optic disc to provide the RNFL thickness of the four quadrants and each of the 12 clock-hour positions. The protocol Macular Cube 512 \times 128 was used to obtain measurements of retinal ganglion cell macular thickness.

All images were obtained by the same experienced technician with a quality score of at least 7/10. Three consecutive scans of the optic disc and macular region were acquired and analysed for each eye. Measurements of RNFL and GCC were averaged using the data of each of the three scans.

2.2. Standard Automated Perimetry. The visual field was assessed by means of a Humphrey Field Analyzer, model 745i II (Carl Zeiss Meditec, Germany), using the 24-2 test program, SITA standard strategy. Near addition was added to the refractive correction, where needed. If fixation losses were greater than 20% and false-positive or false-negative results were higher than 15%, the test was repeated. At least 2 reliable SAPs were performed to minimize the learning effect [35]. Visual field defects were defined as being typically glaucomatous when a standard deviation of the model (PSD) significantly higher than the 5% level and/or a glaucoma hemifield test outside normal limits was recorded.




FIGURE 1: Example of steady-state PERGs of left eyes, presented in time domain (line chart) and frequency domain (bar chart). At a frequency of the stimulus of 7.5 Hz the second harmonic is observed at 15 Hz.

2.3. Pattern Electretinogram. PERG was recorded by means of an instrument supplied by our laboratory (RETIMAX Advanced version 4.3 CSO, Pisa, Italy), using a method similar to the paradigm PERGLA [20], with some minor changes made by our laboratory.

We used as a stimulus horizontal bars with a spatial frequency of 1.7 cycles/degree, which resulted from previous studies as the most sensitive in detecting RGCs dysfunction in early glaucoma [36, 37], modulated in counterphase at 15 reversals/second and electronically generated on a high-resolution LCD monitor (contrast: 90%; luminance: 80 cd/m²; field size: 24° [width] × 24° [height]). The subjects had undilated pupils, of size between 3 and 4 mm, with an appropriate correction for the working distance (57 cm). The signals were recorded from a skin electrode 9 mm Ag/AgCl placed on the lower eyelid. A similar electrode, placed on the lid of the not stimulated eye, was used as a reference, as described in other studies [38].

In all cases the impedance was below 5 k. The responses were amplified (gain of 100.000), filtered (bandwidth: 130 Hz), and sampled with a resolution of 12 bits. The analysis time was 133 ms, equal to the time of presentation of the stimulus (Figure 1). An average (100 events), with automatic rejection of artefacts, was obtained. Five consecutive tests were recorded with a short break, so the duration of the examination was no more than 5 minutes per eye (the total duration being no longer than a visual field examination). The data were then exported to a text file. The amplitude (µV) and phase (π rad) of the second harmonic were then analysed with the Fourier transform (Figures 2 and 3) using a special software programmed by one of the authors (Alberto Mavilio).

The repeatability of the amplitude and phase of the second harmonic was calculated as coefficients of variation (CV, the ratio of the measurement standard deviation to the mean), CVamp (coefficient of variation of amplitude), and CVphase (coefficient of variation of phase), respectively, and as the intraclass correlation coefficients (ICC, describing proportion of total variance accounted for by within-subject variation).




FIGURE 2: Example of steady-state PERGs recorded in response to a pattern of horizontal gratings (1.7 cy/deg. 90% contrast; 80 cd/m² mean luminance; field size: 24° [width] × 24° [height]) alternating at 15 times/s or every 66.6 ms.

The noise level obtained by recording a response to an occluded stimulus was $\leq 0.097 \pm 0.04 \mu V$ in both normal subjects and patients.

In our laboratory the phase decreases with increasing peak time; that is, a delayed response corresponds to lower phase values.

At a reversal rate of 15 Hz the modulo value of 2π rad corresponds to 66.6 ms ($1/15 * 1000 = 66.6$ ms). As described previously [31], to avoid the inherent discontinuity of phase, the recorded value was subtracted from the value of the modulo (2 less than the recorded value). This is needed to prevent negative values of the phase that may affect the calculation of the coefficient of variation.

Statistical analyses were performed using a commercially available software (MedCalc 13.3.1.0). A *P* value of ≤ 0.05 was considered statistically significant.

3. Results

Demographic, structural, and functional data are shown in Tables 1, 2, and 3.

The differences of the variables between groups were analysed using one-way ANOVA analysis of variance, with

TABLE 1: Demographic data.

	Patients						P value*
	EG (37)		GS (21)		NC (24)		
	Mean	SD \pm	Mean	SD \pm	Mean	SD \pm	
Age	57.1	11.6	56.1	10.5	53	6.3	$P = 0.46$
Male (%)	40.5		42.4		41.5		$P = 0.078^{**}$
IOP (mmHg)	15.6	1.1	17.9 ^{ac}	1.6	15.1	1.7	$P < 0.001$
CCT (μm)	546.6	26.6	553.1	32.1	558.7	19.1	$P = 0.337$
MD (dB)	-3.010 ^{ab}	1.95	0.004	1.04	-0.02	1.5	$P < 0.001$
PSD (dB)	3.45 ^{ab}	2.04	1.59	0.49	1.47	0.23	$P < 0.001$
RNFL (μm)	81.56 ^{ab}	9.26	90.8	7.5	94.9	10.1	$P < 0.001$
GCC (μm)	76.13 ^{ab}	6.88	82.09	6.53	84.7	4.9	$P < 0.001$
Amplitude (μV)	0.96 ^{ab}	0.33	0.96 ^a	0.27	1.2	0.26	$P = 0.028$
Phase (π rad)	-0.06	0.36	-0.01	0.26	0.19	0.38	$P = 0.069$
CV_amp (%)	20.25 ^{ab}	13.52	14.27 ^a	7.05	9.42	3.99	$P = 0.004$
CV_phase (%)	8.97 ^b	2.52	7.30 ^a	2.51	3.4	1.13	$P < 0.001$

*One-way analysis of variance (Bonferroni corrected).

**Chi-square.

^aStatistically significant difference from normal control (NC).

^bStatistically significant difference from glaucoma suspects (GS).

^cStatistically significant difference from early manifest glaucoma patients (EG).




FIGURE 3: Fourier spectrum of 5 consecutive tests of steady-state PERG in the same subject. The bar chart shows the amplitude expressed in μV of second harmonic of the signal in response to stimulus of 7.5 Hz.

Bonferroni adjustment. Both linear and logarithmic regressions were used for structural measurements; RNFL and GCC were expressed in linear units (micrometers); functional measurements, PSD, and MD were expressed in logarithmic units (decibels). Pearson correlation in PERG amplitude, phase, coefficients of variation, thickness measurements SD-OCT, and visual field indices are shown in Table 4.

Age, which can affect the amplitude of the PERG and retinal thickness, was not a confounding factor; in fact there was no statistically significant difference of age between all groups (see Table 1).

Because the RNFL thickness shows a slight decrease with age, as suggested by previous studies [39, 40], the data were adjusted for age by a factor of $0.2 \mu\text{m}/\text{year}$ ($0.18\%/ \text{year}$).

The IOP values were significantly higher in the GS group ($17.95 \pm 1.65 \text{ mmHg}$) than in the EG group ($15.67 \pm 1.13 \text{ mmHg}$), because the EG patients were in drug treatment.

PERG amplitude was reduced in EG ($0.96 \pm 0.33 \mu\text{V}$) and GS ($0.96 \pm 0.27 \mu\text{V}$) subjects with respect to NC ($1.20 \pm 0.26 \mu\text{V}$, $P < 0.0001$) and was weakly associated with RNFL thickness ($r = 0.444$, $P = 0.0059$).

CVamp correlates negatively with GCC ($r = -0.379$, $P = 0.0206$), while CVphase correlates better with RNFL ($r = 0.427$, $P = 0.0083$) than with CGG ($r = 0.361$, $P = 0.0283$).

Finally, CVphase correlates fairly strongly with PSD ($r = -0.524$, $P < 0.0009$) and weakly with IOP ($r = 0.362$, $P = 0.0277$).

When the data from normal eyes were entered into a multiple linear regression analysis, with CVphase and CVamp as independent variables and PSD as dependent variable, the significance of the coefficient of variation for CVphase term and for CVamp term was not significant.

In a similar analysis of the EG group, the significance of the coefficient for the CVphase term was $P = 0.0010$ and for the CVamp term was $P = 0.9767$. Table 5 shows the intraclass correlation coefficients (ICC).

4. Discussion

Our results are similar to previous studies. Many authors evaluated the PERG procedure which inspired us in this trial, the PERGLA paradigm, with regard to its reliability in the early diagnosis of glaucoma [16, 18, 19, 26, 27, 41–43].

Several studies also verified structure-function relationship using PERGLA paradigm and methods of analysis of the structure as retinal OCT [13, 28, 41, 44, 45].

Other studies reported that the total thickness of the retina is a good surrogate for glaucomatous damage ganglion

TABLE 2: Clinical characteristics of 37 early glaucoma patients.

EGp	Age	Eye	ph1 (π rad)	ph2 (π rad)	ph3 (π rad)	ph4 (π rad)	ph5 (π rad)	amp1 (μ V)	amp2 (μ V)	amp3 (μ V)	amp4 (μ V)	amp5 (μ V)	MD (db)	PSD (db)	RNFL (μ m)	GCC (μ m)	CV_p (%)	CV_a (%)	Phase (π rad)	Ampl (μ V)	SD_p	SD_a	CCT (μ m)	IOP (mmHg)
#1	49	R	0.09	0.40	0.16	0.33	-0.11	1.57	1.47	1.44	1.43	1.38	-3	2.84	89	83	8.32	4.18	0.17	1.46	0.18	0.06	560	16
#2	49	L	0.27	0.44	0.72	0.60	0.33	1.52	1.24	0.81	0.32	1.37	-2	7	75	70	6.67	41.5	0.47	1.05	0.17	0.44	543	15
#3	69	R	0.24	0.35	0.18	0.19	-0.01	0.73	0.60	0.63	0.75	0.57	-1.02	1.89	82	80	5.42	10.8	0.19	0.66	0.12	0.07	515	14
#4	62	R	0.12	0.42	0.52	0.63	0.63	1.61	1.87	1.70	1.86	1.80	-4	4.29	92	83	7.76	5.52	0.46	1.77	0.19	0.1	537	15
#5	46	L	-0.61	-0.43	-0.80	-0.42	-0.30	1.11	0.81	0.84	0.95	0.79	-0.74	1.25	82	78	11.3	13.5	-0.53	0.9	0.17	0.12	595	17
#6	65	R	-0.44	-0.30	-0.58	-0.42	-0.29	1.34	0.72	1.11	1.06	0.74	-5.9	7.85	74	71	6.53	23.6	-0.41	0.99	0.1	0.23	518	14
#7	60	L	-0.16	0.34	-0.03	0.09	-0.04	1.20	1.04	1.12	0.96	1.01	-2.14	4.36	63	77	8.39	7.93	0.04	1.07	0.17	0.08	540	14
#8	45	R	-0.49	-0.80	-0.55	-0.45	-0.77	0.70	0.60	0.18	0.40	0.91	-6.25	1.76	95	82	10.5	44.9	-0.61	0.56	0.15	0.25	612	17
#9	63	L	0.01	-0.44	0.20	0.32	-0.29	1.03	0.70	0.94	0.88	0.71	-3.54	2.65	100	94	14.6	15.1	-0.04	0.85	0.29	0.13	560	15
#10	35	R	0.22	-0.18	0.12	0.27	0.41	1.85	1.45	1.29	1.46	1.30	-6.11	2.18	71	77	9.06	13.7	0.17	1.47	0.2	0.2	550	16
#11	59	L	-0.26	0.32	0.47	0.06	0.35	1.11	0.67	0.78	0.76	0.79	-2.01	1.22	73	70	11.9	18.4	0.19	0.82	0.26	0.15	530	15
#12	39	L	-0.37	-0.10	-0.19	0.27	0.10	1.21	0.77	1.04	0.79	0.82	-1.5	1.87	83	77	11.5	18.3	-0.06	0.93	0.22	0.17	525	15
#13	57	R	0.13	-0.08	0.14	-0.01	0.20	1.42	1.48	1.48	1.11	1.00	-5.11	7.75	77	67	5.07	18.5	0.08	1.19	0.11	0.22	570	16
#14	54	L	-0.51	-0.31	-0.22	0.01	0.07	1.37	1.17	1.16	1.35	0.97	-3.21	2.79	97	85	11.7	12.4	-0.19	1.2	0.21	0.15	595	17
#15	59	L	0.36	0.76	0.10	0.27	0.36	1.05	0.79	1.10	1.03	1.11	-1.11	2.94	95	84	9.1	11.6	0.37	1.02	0.22	0.12	560	16
#16	75	R	-0.72	-0.08	-0.20	-0.06	-0.10	0.76	1.02	0.75	0.82	0.83	-2.13	3.55	87	88	14.1	74.6	-0.23	0.82	0.25	0.06	525	17
#17	65	R	-0.31	-0.59	-0.29	-0.31	-0.46	0.77	1.03	1.05	0.60	0.87	-3.57	2.61	92	86	7.09	19.4	-0.39	0.87	0.11	0.17	505	16
#18	60	R	0.19	0.32	-0.08	0.19	-0.19	0.56	0.52	0.87	0.56	0.62	-5.65	3.96	85	69	9.07	20	0.09	0.63	0.19	0.12	543	15
#19	81	R	-0.97	-0.63	-0.86	-0.71	-0.92	1.13	0.40	0.79	0.39	1.01	0.35	1.38	83	68	10.8	40.9	-0.82	0.75	0.13	0.3	575	16
#20	45	R	0.22	0.58	0.51	0.66	0.33	1.41	1.52	1.49	0.92	1.07	-1.2	6.3	73	69	6.58	18.9	0.46	1.28	0.16	0.24	555	15
#21	40	L	-0.11	0.11	0.32	0.18	0.55	1.82	1.58	1.06	1.85	1.46	-2.06	2.38	77	74	10	18.6	0.21	1.55	0.22	0.29	523	14
#22	72	L	0.48	0.12	0.50	0.37	0.48	1.18	0.93	1.19	0.98	0.83	-2.89	3.03	78	72	5.88	13.8	0.39	1.02	0.14	0.14	545	14
#23	52	R	-0.57	-0.38	-0.04	-0.21	-0.31	1.45	1.54	1.39	1.13	1.55	-4.5	1.62	90	82	10.3	10.8	-0.3	1.41	0.17	0.15	569	15
#24	76	L	0.12	-0.28	0.03	-0.07	-0.44	0.35	1.59	0.66	0.35	0.35	-6.49	3.76	78	66	10.9	72.7	-0.13	0.66	0.2	0.48	555	15
#25	40	R	-0.20	0.08	-0.06	0.03	0.09	1.02	0.64	0.75	0.71	0.79	-3.22	5.72	69	77	5.55	16.8	-0.01	0.78	0.11	0.13	525	16
#26	49	R	0.04	-0.04	0.07	0.27	-0.41	1.16	1.03	0.99	1.00	1.18	-2.15	1.08	77	67	11.3	7.81	-0.02	1.07	0.22	0.08	495	17
#27	43	L	-0.37	-0.19	0.05	-0.14	-0.21	0.88	1.67	1.28	1.21	0.71	-1.52	2.33	88	74	7.5	29.1	-0.17	1.15	0.14	0.33	536	15
#28	68	L	-0.09	-0.41	-0.04	-0.49	-0.07	1.18	1.38	0.88	1.22	1.04	-0.06	1.48	96	84	10.7	15	-0.22	1.14	0.19	0.17	520	16
#29	49	R	0.75	0.14	0.57	0.58	0.54	0.88	0.85	0.64	1.04	0.57	-4.55	2.24	71	74	7.99	21.2	0.52	0.8	0.2	0.17	578	17
#30	70	R	0.45	0.68	0.22	0.74	0.26	0.61	0.61	0.52	0.51	0.38	-0.45	1.59	78	76	8.64	15.5	0.47	0.53	0.21	0.08	542	17
#31	67	R	-0.43	-0.82	-0.45	-0.45	-0.45	0.40	0.34	0.29	0.40	0.77	-5.94	7.15	80	78	11.6	38.5	-0.49	0.44	0.18	0.17	523	17
#32	42	R	-0.71	-0.91	-0.86	-0.62	-0.72	0.24	0.26	0.54	0.44	0.63	-2.26	3.45	86	69	8.52	36	-0.76	0.42	0.11	0.15	555	18
#33	65	R	-0.41	-0.17	0.04	0.15	-0.05	1.64	1.17	1.02	1.31	1.02	-2.25	3.32	88	73	10.1	18.7	-0.09	1.23	0.19	0.23	567	17
#34	62	L	-0.85	-0.77	-0.82	-0.71	-0.49	0.43	0.29	0.48	0.42	0.24	-1.16	1.69	76	79	10.1	24.7	-0.73	0.37	0.13	0.09	545	17
#35	63	L	-0.26	-0.10	-0.22	0.02	-0.24	1.38	1.32	1.11	1.43	1.24	-2.06	3.92	81	72	5.7	8.65	-0.16	1.3	0.11	0.11	515	15
#36	58	R	-0.04	-0.01	0.16	0.06	-0.03	1.12	1.18	0.81	1.05	0.67	-6.71	7.78	60	76	3.74	19.9	0.03	0.97	0.08	0.19	535	15
#37	61	L	-0.33	-0.29	0.07	-0.20	-0.26	0.67	0.75	0.56	0.48	0.65	-3.25	5.02	70	76	8.09	15.3	-0.2	0.62	0.15	0.1	585	14

PERG phase and amplitude (values of 5 consecutive tests), mean deviation (MD), pattern standard deviation (PSD), retinal nerve fiber layer thickness (RNFL), ganglion cell complex (GCC), coefficient of variation of amplitude (CV_a), coefficient of variation of phase (CV_p), mean phase (phase), mean amplitude (Ampl), standard deviation of amplitude (SD_a), standard deviation of phase (CV_p), central corneal thickness (CCT), and intraocular pressure (IOP) in 37 early glaucoma patients (EGp).

TABLE 3: Clinical characteristics of 21 glaucoma suspect patients.

SGp	Age	Eye	ph1 (π rad)	ph2 (π rad)	ph3 (π rad)	ph4 (π rad)	ph5 (π rad)	amp1 (μ V)	amp2 (μ V)	amp3 (μ V)	amp4 (μ V)	amp5 (μ V)	MD (db)	PSD (db)	RNFL (μ m)	GCC (μ m)	CV_p (%)	CV_a (%)	Phase (π rad)	Ampl (μ V)	SD_p	SD_a	CCT (μ m)	IOP (mmHg)
#1	47	R	0.60	0.20	0.25	0.33	0.16	1.35	1.30	1.32	1.49	1.13	1.51	1.36	86	83	6.79	8.75	0.31	1.32	0.16	0.12	535	16
#2	54	R	0.04	-0.30	-0.37	-0.20	-0.08	1.00	1.01	0.88	0.83	0.76	-0.23	1.24	92	90	8.09	10.67	-0.18	0.9	0.15	0.1	570	16
#3	49	L	0.12	-0.24	-0.01	0.17	0.10	1.10	1.08	0.95	0.95	0.92	-0.59	1.78	90	78	7.23	7.69	0.03	1	0.15	0.08	597	18
#4	53	R	-0.20	0.12	0.04	-0.04	-0.08	0.95	0.73	0.87	0.89	0.70	-0.21	1.88	91	86	5.52	11.72	-0.03	1	0.11	0.1	515	18
#5	58	L	0.33	0.24	0.01	0.32	0.21	1.47	1.35	1.09	1.51	1.19	-0.78	1.88	84	79	5.11	12.02	0.22	1.32	0.11	0.16	565	20
#6	57	R	-0.16	0.17	0.27	-0.30	0.10	0.59	0.43	0.48	0.66	0.49	0.62	1.16	84	98	10.5	15.79	0.02	0.53	0.21	0.08	525	18
#7	73	L	-0.06	-0.03	-0.04	-0.08	0.17	1.43	1.05	0.87	1.08	1.23	-0.33	1.63	93	74	4.61	16.73	-0.01	1.13	0.09	0.19	549	16
#8	35	R	0.25	0.71	0.66	0.79	0.74	1.58	1.16	1.29	1.62	1.31	-0.25	1.23	96	78	7.3	12.76	0.63	1.39	0.19	0.18	508	16
#9	63	R	0.04	0.42	0.14	0.07	0.14	0.89	0.82	0.98	0.64	0.89	-0.74	1.87	108	91	6.24	13.44	0.16	1.01	0.13	0.11	578	16
#10	77	R	0.14	0.00	0.15	0.01	0.24	0.97	0.56	0.98	0.67	0.64	0.65	1.24	93	72	4.31	23.01	0.11	0.92	0.09	0.18	562	17
#11	53	L	0.42	0.33	0.11	0.37	0.38	0.94	0.65	0.69	0.54	0.66	0.02	1.33	80	80	4.72	19.1	0.32	0.7	0.11	0.13	601	18
#12	64	L	-0.04	0.00	-0.39	-0.16	-0.11	0.70	0.89	0.34	0.37	0.40	-0.01	1.23	96	88	7.25	40.32	-0.14	0.65	0.13	0.22	555	17
#13	50	L	-0.52	-0.64	-0.13	-0.34	-0.33	1.00	0.58	0.65	0.78	0.89	-0.66	1.33	87	82	10.78	19.59	-0.39	0.94	0.17	0.15	495	22
#14	62	L	-0.02	0.20	-0.03	0.23	0.14	0.53	0.40	0.51	0.59	0.48	0.68	1.1	75	74	5.25	12.25	0.1	0.5	0.11	0.06	575	18
#15	57	R	-0.51	-0.68	-0.66	-0.33	-0.62	1.34	1.54	0.86	1.03	0.65	0.84	1.24	93	86	8.93	29.8	-0.56	1.08	0.13	0.32	535	18
#16	64	L	-0.31	-0.15	-0.05	-0.07	0.20	0.66	0.67	0.73	0.55	0.72	1.43	1.4	87	73	8.52	9.33	-0.07	0.8	0.16	0.06	560	21
#17	59	L	-0.12	-0.02	0.05	-0.16	0.01	1.29	1.56	1.45	1.16	1.60	0.27	1.53	94	83	4.04	11.75	-0.05	1.41	0.08	0.17	525	18
#18	51	L	-0.41	0.00	-0.33	-0.12	-0.40	0.65	0.64	0.66	0.66	0.58	0.88	1.43	93	84	9.36	5.09	-0.25	0.77	0.16	0.03	565	18
#19	32	L	-0.09	0.18	0.12	0.03	0.21	1.40	1.42	1.12	1.17	1.24	-0.37	1.34	89	82	5.29	9.76	0.09	1.27	0.11	0.12	608	18
#20	60	L	-0.30	-0.46	-0.22	-0.52	0.13	0.78	0.70	0.66	0.47	0.85	0.25	2.08	106	85	13.18	18.58	-0.27	0.69	0.23	0.13	512	18
#21	61	R	-0.22	-0.17	0.00	-0.35	-0.54	0.82	0.64	0.63	0.88	0.88	0.99	1.45	90	78	10.38	14.46	-0.26	0.93	0.18	0.11	582	20

PERG phase and amplitude (values of 5 consecutive tests), mean deviation (MD), pattern standard deviation (PSD), retinal nerve fiber layer thickness (RNFL), ganglion cell complex (GCC), coefficient of variation of amplitude (CV_a), coefficient of variation of phase (CV_p), mean phase (phase), mean amplitude (Ampl), standard deviation of amplitude (SD_a), standard deviation of phase (CV_p), central corneal thickness (CCT), and intraocular pressure (IOP) in 21 glaucoma suspect patients (SGp).

TABLE 4: Pearson correlation coefficient (CC) and significance level P (SL- P) in 37 early manifest glaucoma patients.

	Age	IOP (mmHg)	CCT (μm)	MD (dB)	PSD (dB)	RNFL (μm)	GCC (μm)	Amplitude (μV)	Phase (π rad)	CV_amp (%)	CV_phase (%)
Age	CC	-0.108	-0.133	0.082	0.038	0.133	0.051	-0.181	-0.128	0.126	0.105
	SL- P	0.523	0.433	0.629	0.823	0.432	0.762	0.284	0.449	0.457	0.536
IOP (mmHg)	CC	-0.108	0.269	0.078	-0.252	0.296	0.177	-0.251	-0.364	0.11	0.362
	SL- P	0.523	0.108	0.646	0.133	0.075	0.294	0.134	0.026	0.516	0.0277
CCT (μm)	CC	-0.133	0.269	-0.131	-0.108	0.181	0.019	0.043	-0.157	0.207	0.178
	SL- P	0.433	0.108	0.441	0.526	0.282	0.908	0.8	0.352	0.218	0.292
MD (dB)	CC	0.082	0.078	-0.131	-0.465	0.117	0.05	0.025	0.011	-0.287	0.153
	SL- P	0.629	0.646	0.441	0.0037	0.491	0.766	0.882	0.949	0.084	0.366
PSD (dB)	CC	0.038	-0.252	-0.108	-0.465	-0.386	-0.279	0.024	0.138	0.152	-0.524
	SL- P	0.823	0.133	0.0037	0.0037	0.018	0.0945	0.889	0.414	0.369	0.0009
RNFL (μm)	CC	0.133	0.296	0.117	-0.386	0.636	0.636	0.444	-0.228	-0.05	0.427
	SL- P	0.432	0.075	0.282	0.0184	<0.0001	<0.0001	0.005	0.175	0.768	0.0083
GCC (μm)	CC	0.051	0.177	0.05	-0.279	0.636	0.636	0.24	-0.054	-0.379	0.361
	SL- P	0.762	0.294	0.766	0.094	<0.0001	<0.0001	0.152	0.753	0.0206	0.0283
Amplitude (μV)	CC	-0.181	0.043	0.025	0.024	0.444	0.24	0.444	0.285	-0.32	-0.037
	SL- P	0.284	0.8	0.882	0.889	0.0059	0.152	0.444	0.087	0.0536	0.828
Phase (π rad)	CC	-0.128	-0.364	0.011	0.138	-0.228	-0.054	0.285	-0.346	-0.346	-0.33
	SL- P	0.449	0.0268	0.352	0.414	0.175	0.753	0.087	0.0358	0.0358	0.0461
CV_amp (%)	CC	0.126	0.11	0.207	0.152	-0.05	-0.379	-0.32	-0.346	0.09	0.09
	SL- P	0.457	0.516	0.218	0.369	0.768	0.02	0.053	0.0358	0.0358	0.597
CV_phase (%)	CC	0.105	0.362	0.178	-0.524	0.427	0.361	-0.037	-0.33	0.09	0.09
	SL- P	0.536	0.0277	0.292	0.0009	0.0083	0.028	0.828	0.0461	0.597	0.597

Intraocular pressure (IOP), central corneal thickness (CCT), mean deviation (MD), pattern standard deviation (PSD), retinal nerve fiber layer thickness (RNFL), ganglion cell complex (GCC), PERG amplitude, PERG phase, coefficient of variation PERG amplitude (CV_amp), and coefficient of variation PERG phase (CV_Phase).

TABLE 5: Intraclass correlation coefficient.

	EG (37)		Patients GS (21)		NC (24)	
	ICC ^a	95% CI	ICC ^a	95% CI	ICC ^a	95% CI
PERG amplitude (μV)	0.9187	0.8684 to 0.9538	0.9466	0.8993 to 0.9756	0.9642	0.9226 to 0.9869
PERG phase (π rad)	0.9459	0.9124 to 0.9692	0.9232	0.8554 to 0.9650	0.9879	0.9739 to 0.9956

^aThe degree of consistency among measurements.

ICC = intraclass correlation coefficient.

IC = confidence interval.

cell layer measured as SD-OCT [46, 47] and that the total macular thickness was significantly associated with glaucoma [7]. Nevertheless, we observed a better association between PSD, which indicates the severity of the disease, and RNFL ($R^2 = 0.1487$, $P = 0.0184$) than GCC ($R^2 = 0.07782$, $P = 0.0945$).

Longitudinal studies have shown that the PERG amplitude is able to detect signs of glaucomatous damage before psychometric and morphological techniques [15, 48, 49].

The PERG amplitude was significantly lower in EG and GS patients ($0.96 \pm 0.33 \mu V$ and $0.96 \pm 0.27 \mu V$, resp.) compared to healthy ($1.20 \pm 0.26 \mu V$). Our data demonstrate that the reduction of PERG amplitude in SG subjects that, unlike EG patients, still do not show abnormal visual field and reduced retinal thickness, is an indicator of RGCs dysfunction. As noted by other authors [13] PERG amplitude is weakly but significantly associated with RNFL thickness reduction in EG patients ($R^2 = 0.1975$, $P = 0.0059$), whereas PERG phase in the current study did not show statistically significant differences between the subjects of the different groups.

It must be said that our trial PERG amplitude was higher than that previously reported by Bowd et al. [18] ($0.96 \mu V$ compared to $0.83 \mu V$) and similar to that previously reported by in healthy eyes ($1.2 \mu V$ compared to $1.1 \mu V$ [20]). This can be explained by the different degree of severity of glaucoma, worse in Bowd patients (MD -9.0 dB), compared to -3.0 dB found in our EG patients.

Amplitude in the first test tended to be greater than that in successive tests. This finding may be related to the percentage of decrease in the amplitude due to adaptation to the stimulus PERG [30]. For this reason we have focused our attention on CVphase.

It is known that amplitude and phase of the PERG represent two distinct aspects of neural activity [31]. Briefly, the amplitude is related to the number of neurons; the phase delay is a further indicator of viability of activated neurons that may or may not be associated with amplitude reduction and in particular may mean that RGCs active respond more slowly. The last becomes progressively delayed with aging [20] and may be further delayed in early glaucoma [21].

Falsini et al. [44] showed that the loss of RGCs electrophysiological function is relatively greater than expected from the anatomical loss of RGCs axons in early glaucoma and that RNFL thickness reduction, less than expected, could be explained by glial remodelling.

In addition, changes in dendrites typically precede neuronal loss and lead to a reduction in the responsiveness of RGCs in glaucoma [50].

PERG generation includes a mixed population of RGCs [51], which are commonly divided into two major classes: P-cells (approximately 80% of the total) and M-cells (approximately 10%). M-cells are much more sensitive to changes than P-cells [52], and their response is temporally faster than P-cells [53], the first responding promptly to the steady-state PERG stimulus.

Since M-cells are relatively few and sparse, even a partial malfunction can be early detected by doubling perimetry frequency [54]. The increase of CVphase in SG subjects suggests that in early glaucoma there would be a progressive loss of the ability of RGCs to adapt in response to the growing demand for energy associated with the high-contrast PERG stimulus [55].

Porciatti and Ventura reported that PERG phase delays suggest pathophysiological mechanisms such as dendritic dysfunction or delay in axonal transport and may represent an opportunity to detect RGCs dysfunction preceding cell death [30].

In this study we demonstrated a significant but modest relationship between the RGCs function using steady-state PERG and some structural measures such as GCC and RNFL measured by OCT (see Table 4). These results suggest that a population of viable RGCs responds with reduced activity, thus signalling a period of discomfort that precedes cell death.

The primary purpose of this study was to evaluate the variability of the PERG signal because, in our opinion, it is increased more in patients suffering from glaucoma than in healthy subjects. In glaucoma suspects, reduced amplitude and increased variability of the phase of the PERG may indicate the presence of a functional impairment.

In fact, the variability of PERG signal has been evaluated in several studies: Bowd et al. [16] studied medium amplitudes and phases, their noise level, SNR, within-subject variability, CV, and ICC PERGLA recordings for within and between trials.

For amplitude, Bowd et al. observed that the variability of successive measurements was approximately 10% to 12% for healthy eyes and glaucoma patients, respectively. These values were similar within and between trials.

As for the phase, repeat measurement variability observed in this study was very low, about 1% to 2%; furthermore, ICC indicated that the percentage of total variance accounted for by intrasubject variation was similar in within and between trials and for healthy eyes and patient (range: 82% to 92%).

We observed within-trial variability of the amplitude higher than previous study, on the order of 15–20% for glaucoma patients and 10% for controls. The variability of




FIGURE 4: Five consecutive tests of steady-state PERG in the same subject. The line chart shows the trend of the phase in the frequency domain from 7.5 Hz to 30 Hz. The phase always passes from the same point at 15 Hz that corresponds to the second harmonic of the signal in response to stimulus of 7.5 Hz.

the phase was also higher than quoted study, of 8–10% for glaucomatous eyes versus 3–4% for healthy subjects.

In this study the variability of the phase has been observed not as an element of repeatability of the method but as a discriminating feature between healthy and diseased eyes. This approach can help to detect nonspecific reduction in the amplitude of PERG.

In fact, the phase is not delayed when the contrast stimulus is artificially deteriorated by simulating the visual acuity reduction due to the cataract [30]; that is, conditions like cataract reduce the PERG amplitude but do not influence the phase delay. In other words, while the PERG amplitude and its variability may depend on the opacity of the media, the phase delay and its variability, as showed in our work, are not affected by the same conditions and may express RGCs dysfunction.

In our study we did not evaluate phase delay, but its greater variability that may be a sign of RGCs dysfunction.

If you look at the phase in the frequency domain (Figure 4), by repeating the test you will see that it has an almost chaotic behavior but passes always from the same point that corresponds with the frequency of the second harmonic.

CVamp and CVphase were significantly lower in healthy individuals ($3.54 \pm 1.13\%$), compared to the groups GS ($7.30 \pm 2.51\%$) and EG ($8.97 \pm 2.52\%$), respectively (see Table 1).

Our work did not include between-proof recordings, but Fredette et al. [26] observed the test-retest variability of the PERG amplitude, expressed in terms of standard deviation (SD) of the results obtained for each subject of glaucoma during the 5 sessions in 5 different days. They also assessed the intrinsic variability (intratest), which was defined as the standard deviation of 2 consecutive recordings divided by the square root of 2. Fredette et al. argue that the amplitude variability could not be used to discriminate healthy from the glaucomatous eyes because they attributed the greater variability amplitude to the reduced amplitude of the signal in glaucomatous patients. In fact, mean amplitude and variability in their work were correlated significantly

($r^2 = 0.164$, $P = 0.003$), indicating that the variability amplitude was due to the low SNR.

Our trial showed no significant association between amplitude and SD ($r^2 = 0.010$, $P = 0.546$), probably because the amplitude of our signal and probably also our SNR were higher than measurement previously reported by Fredette et al. ($0.09 \pm 0.04 \mu\text{V}$ versus $0.08 \pm 0.03 \mu\text{V}$).

In addition, our trial showed no association between PERG amplitude and CVphase ($r^2 = 0.001$, $P = 0.821$), confirming that variability of the phase was not dependent on the magnitude of the signal.

The structure-function relationship between CVphase and RNFL or GCC was significant only in EG group (Table 4).

The intraclass correlation coefficients indicated that the percentage of the total variance represented by the within-trial variation (i.e., the measurement reliability) was very low with regard to both the phase (0.9124 to 0.9692 95% CI) and the amplitude (0.8684 to 0.9538 95% CI), so our procedure appeared reliable.

5. Conclusions

To our knowledge, this is the first approach focusing on the intrasession phase variability. The evaluation of intraindividual and intrasession variability signal has undeniable advantages in that it minimizes interindividual variations.

This research has several limitations. In our study, we limited the variables that could contribute to measurement artifacts including cataracts or other media opacities, poor visual acuity, and pupillary miosis, but the loss of fixation of the patient cannot be quantified with our technology.

The test-retest variability should vary with the dynamic range of each instrument, so each laboratory should determine its own variability on a normative sample of healthy subjects. In our study the variability of the signal does not correlate with age, but our control sample is too small to know whether any normative database should be corrected for age.

In clinical practice it is not easy to identify with certainty the actual visual acuity at a distance of work and the transparency of the dioptric media that can compromise the signal-to-noise ratio.

The values of the impedances should be constantly controlled by the operator, but most of the tools do not have these characteristics.

Standardizing the SNR is not an easy task and the assumptions of this work are governed by the reliability and experience of the laboratory, especially when working with a steady-state stimulus [25].

Further studies are needed, but our hypothesis is that increased phase variability of intrasession PERG may represent an opportunity to detect RGCs dysfunction preceding cell death.

Conflict of Interests

None of the authors has conflict of interests with the submission. No financial support was received for this submission.

Acknowledgment

The authors thank Dario Sisto, MD, who gave them kind suggestions and some important fixes.

References

- [1] R. S. Harwerth, L. Carter-Dawson, F. Shen, E. L. Smith III, and M. L. J. Crawford, "Ganglion cell losses underlying visual field defects from experimental glaucoma," *Investigative Ophthalmology and Visual Science*, vol. 40, no. 10, pp. 2242–2250, 1999.
- [2] L. A. Kerrigan-Baumrind, H. A. Quigley, M. E. Pease, D. F. Kerrigan, and R. S. Mitchell, "Number of ganglion cells in glaucoma eyes compared with threshold visual field tests in the same persons," *Investigative Ophthalmology and Visual Science*, vol. 41, no. 3, pp. 741–748, 2000.
- [3] G. Wollstein, H. Ishikawa, J. Wang, S. A. Beaton, and J. S. Schuman, "Comparison of three optical coherence tomography scanning areas for detection of glaucomatous damage," *The American Journal of Ophthalmology*, vol. 139, no. 1, pp. 39–43, 2005.
- [4] G. Wollstein, J. S. Schuman, L. L. Price et al., "Optical coherence tomography (OCT) macular and peripapillary retinal nerve fiber layer measurements and automated visual fields," *The American Journal of Ophthalmology*, vol. 138, no. 2, pp. 218–225, 2004.
- [5] J. L. Hougaard, A. Heijl, and B. Bengtsson, "Glaucoma detection by stratus OCT," *Journal of Glaucoma*, vol. 16, no. 3, pp. 302–306, 2007.
- [6] O. Tan, V. Chopra, A. T.-H. Lu et al., "Detection of macular ganglion cell loss in glaucoma by Fourier-domain optical coherence tomography," *Ophthalmology*, vol. 116, no. 12, pp. 2305–14.e1-2, 2009.
- [7] H. L. Rao, L. M. Zangwill, R. N. Weinreb, P. A. Sample, L. M. Alencar, and F. A. Medeiros, "Comparison of different spectral domain optical coherence tomography scanning areas for glaucoma diagnosis," *Ophthalmology*, vol. 117, no. 9, pp. 1692–1699, 2010.
- [8] S. Mori, M. Hangai, A. Sakamoto, and N. Yoshimura, "Spectral-domain optical coherence tomography measurement of macular volume for diagnosing glaucoma," *Journal of Glaucoma*, vol. 19, no. 8, pp. 528–534, 2010.
- [9] J. Kotowski, G. Wollstein, L. S. Folio, H. Ishikawa, and J. S. Schuman, "Clinical use of OCT in assessing glaucoma progression," *Ophthalmic Surgery, Lasers & Imaging*, vol. 42, pp. S6–S14, 2011.
- [10] C. A. Curcio and K. A. Allen, "Topography of ganglion cells in human retina," *Journal of Comparative Neurology*, vol. 300, no. 1, pp. 5–25, 1990.
- [11] L. Maffei and A. Fiorentini, "Electroretinographic responses to alternating gratings before and after section of the optic nerve," *Science*, vol. 211, no. 4485, pp. 953–955, 1981.
- [12] E. Zrenner, "The physiological basis of the pattern electroretinogram," in *Progress in Retinal Research*, pp. 427–464, Pergamon Press, Oxford, UK, 1990.
- [13] L. M. Ventura, N. Sorokac, R. de Los Santos, W. J. Feuer, and V. Porciatti, "The relationship between retinal ganglion cell function and retinal nerve fiber thickness in early glaucoma," *Investigative Ophthalmology and Visual Science*, vol. 47, no. 9, pp. 3904–3911, 2006.
- [14] M. Bach and M. B. Hoffmann, "Update on the pattern electroretinogram in glaucoma," *Optometry and Vision Science*, vol. 85, no. 6, pp. 386–395, 2008.
- [15] N. Pfeiffer and M. Bach, "The pattern-electroretinogram in glaucoma and ocular hypertension. A cross-sectional and longitudinal study," *German Journal of Ophthalmology*, vol. 1, no. 1, pp. 35–40, 1992.
- [16] C. Bowd, A. Tafreshi, G. Vizzeri, L. M. Zangwill, P. A. Sample, and R. N. Weinreb, "Repeatability of pattern electroretinogram measurements using a new paradigm optimized for glaucoma detection," *Journal of Glaucoma*, vol. 18, no. 6, pp. 437–442, 2009.
- [17] C. Bowd, A. Tafreshi, L. M. Zangwill, F. A. Medeiros, P. A. Sample, and R. N. Weinreb, "Pattern electroretinogram association with spectral domain-OCT structural measurements in glaucoma," *Eye*, vol. 25, no. 2, pp. 224–232, 2011.
- [18] C. Bowd, G. Vizzeri, A. Tafreshi, L. M. Zangwill, P. A. Sample, and R. N. Weinreb, "Diagnostic accuracy of pattern electroretinogram optimized for glaucoma detection," *Ophthalmology*, vol. 116, no. 3, pp. 437–443, 2009.
- [19] A. Tafreshi, L. Racette, R. N. Weinreb et al., "Pattern electroretinogram and psychophysical tests of visual function for discriminating between healthy and glaucoma eyes," *The American Journal of Ophthalmology*, vol. 149, no. 3, pp. 488–495, 2010.
- [20] V. Porciatti and L. M. Ventura, "Normative data for a user-friendly paradigm for pattern electroretinogram recording," *Ophthalmology*, vol. 111, no. 1, pp. 161–168, 2004.
- [21] L. M. Ventura and V. Porciatti, "Restoration of retinal ganglion cell function in early glaucoma after intraocular pressure reduction: a pilot study," *Ophthalmology*, vol. 112, no. 1, pp. 20–27, 2005.
- [22] M. Bach, M. G. Brigell, M. Hawlina et al., "ISCEV standard for clinical pattern electroretinography (PERG): 2012 update," *Documenta Ophthalmologica*, vol. 126, no. 1, pp. 1–7, 2013.
- [23] J. W. Cooley and J. W. Tukey, "An algorithm for the machine calculation of complex fourier series," *Mathematics of Computation*, vol. 19, no. 90, pp. 297–301, 1965.
- [24] M. A. O'Neill, "Faster than fast Fourier," *Byte*, vol. 4, pp. 293–300, 1988.
- [25] M. Bach and T. Meigen, "Do's and don'ts in Fourier analysis of steady-state potentials," *Documenta Ophthalmologica*, vol. 99, no. 1, pp. 69–82, 1999.
- [26] M.-J. Fredette, D. R. Anderson, V. Porciatti, and W. Feuer, "Reproducibility of pattern electroretinogram in glaucoma patients with a range of severity of disease with the new glaucoma paradigm," *Ophthalmology*, vol. 115, no. 6, pp. 957–963, 2008.
- [27] A. Yang and W. H. Swanson, "A new pattern electroretinogram paradigm evaluated in terms of user friendliness and agreement with perimetry," *Ophthalmology*, vol. 114, no. 4, pp. 671–679, 2007.
- [28] M. Araie, "Test-retest variability in structural parameters measured with glaucoma imaging devices," *Japanese Journal of Ophthalmology*, vol. 57, no. 1, pp. 1–24, 2013.
- [29] V. Porciatti, N. Sorokac, and W. Buchser, "Habituation of retinal ganglion cell activity in response to steady state pattern visual stimuli in normal subjects," *Investigative Ophthalmology and Visual Science*, vol. 46, no. 4, pp. 1296–1302, 2005.
- [30] V. Porciatti and L. M. Ventura, "Adaptive changes of inner retina function in response to sustained pattern stimulation," *Vision Research*, vol. 49, no. 5, pp. 505–513, 2009.

- [31] V. Porciatti and L. M. Ventura, "Physiologic significance of steady-state pattern electroretinogram losses in glaucoma: clues from simulation of abnormalities in normal subjects," *Journal of Glaucoma*, vol. 18, no. 7, pp. 535–542, 2009.
- [32] B. L. Lam, W. J. Feuer, F. Abukhalil, V. Porciatti, W. W. Hauswirth, and J. Guy, "Leber hereditary optic neuropathy gene therapy clinical trial recruitment: year 1," *Archives of Ophthalmology*, vol. 128, no. 9, pp. 1129–1135, 2010.
- [33] L. M. Ventura, I. Golubev, W. J. Feuer, and V. Porciatti, "The PERG in diabetic glaucoma suspects with no evidence of retinopathy," *Journal of Glaucoma*, vol. 19, no. 4, pp. 243–247, 2010.
- [34] I. Bodis-Wollner, "Visual electrophysiology in Parkinson's disease: PERG, VEP and visual P300," *Clinical EEG Electroencephalography*, vol. 28, no. 3, pp. 143–147, 1997.
- [35] A. Heijl, G. Lindgren, and J. Olsson, "The effect of perimetric experience in normal subjects," *Archives of Ophthalmology*, vol. 107, no. 1, pp. 81–86, 1989.
- [36] V. Porciatti, B. Falsini, G. Scalia, A. Fadda, and G. Fontanesi, "The pattern electroretinogram by skin electrodes: effect of spatial frequency and age," *Documenta Ophthalmologica*, vol. 70, no. 1, pp. 117–122, 1988.
- [37] L. M. Ventura, V. Porciatti, K. Ishida, W. J. Feuer, and R. K. Parrish II, "Pattern electroretinogram abnormality and glaucoma," *Ophthalmology*, vol. 112, no. 1, pp. 10–19, 2005.
- [38] T. Salgarello, A. Colotto, B. Falsini et al., "Correlation of pattern electroretinogram with optic disc cup shape in ocular hypertension," *Investigative Ophthalmology and Visual Science*, vol. 40, no. 9, pp. 1989–1997, 1999.
- [39] A. Kanamori, M. Nakamura, M. Tomioka, Y. Kawaka, Y. Yamada, and A. Negi, "Structure-function relationship among three types of spectral-domain optical coherent tomography instruments in measuring parapapillary retinal nerve fiber layer thickness," *Acta Ophthalmologica*, vol. 91, no. 3, pp. e196–e202, 2013.
- [40] P. Sony, R. Sihota, H. K. Tewari, P. Venkatesh, and R. Singh, "Quantification of the retinal nerve fiber layer thickness in normal Indian eyes with optical coherence tomography," *Indian Journal of Ophthalmology*, vol. 52, no. 4, pp. 303–309, 2004.
- [41] M. Sehi, M. Pinzon-Plazas, W. J. Feuer, and D. S. Greenfield, "Relationship between pattern electroretinogram, standard automated perimetry, and optic nerve structural assessments," *Journal of Glaucoma*, vol. 18, no. 8, pp. 608–617, 2009.
- [42] R. Forte, L. Ambrosio, P. Bonavolontá, and G. Ambrosio, "Pattern electroretinogram optimized for glaucoma screening (PERGLA) and retinal nerve fiber thickness in suspected glaucoma and ocular hypertension," *Documenta Ophthalmologica*, vol. 120, no. 2, pp. 187–192, 2010.
- [43] G. Vizzeri, A. Tafreshi, R. N. Weinreb, and C. Bowd, "Effect of operator and optical defocus on the variability of pattern electroretinogram optimized for glaucoma detection (PERGLA)," *Journal of Glaucoma*, vol. 19, no. 2, pp. 77–82, 2010.
- [44] B. Falsini, D. Marangoni, T. Salgarello et al., "Structure—function relationship in ocular hypertension and glaucoma: interindividual and interocular analysis by OCT and pattern ERG," *Graefes' Archive for Clinical and Experimental Ophthalmology*, vol. 246, no. 8, pp. 1153–1162, 2008.
- [45] M. R. Banitt, L. M. Ventura, W. J. Feuer et al., "Progressive loss of retinal ganglion cell function precedes structural loss by several years in glaucoma suspects," *Investigative Ophthalmology and Visual Science*, vol. 54, no. 3, pp. 2346–2352, 2013.
- [46] V. Guedes, J. S. Schuman, E. Hertzmark et al., "Optical coherence tomography measurement of macular and nerve fiber layer thickness in normal and glaucomatous human eyes," *Ophthalmology*, vol. 110, no. 1, pp. 177–189, 2003.
- [47] T. Ojima, T. Tanabe, M. Hangai, S. Yu, S. Morishita, and N. Yoshimura, "Measurement of retinal nerve fiber layer thickness and macular volume for glaucoma detection using optical coherence tomography," *Japanese Journal of Ophthalmology*, vol. 51, no. 3, pp. 197–203, 2007.
- [48] S. F. N. Bode, T. Jehle, and M. Bach, "Pattern electroretinogram in glaucoma suspects: new findings from a longitudinal study," *Investigative Ophthalmology and Visual Science*, vol. 52, no. 7, pp. 4300–4306, 2011.
- [49] M. Bach, A. S. Unsoeld, H. Philipppin et al., "Pattern ERG as an early glaucoma indicator in ocular hypertension: a long-term, prospective study," *Investigative Ophthalmology & Visual Science*, vol. 47, no. 11, pp. 4881–4887, 2006.
- [50] A. J. Weber and C. D. Harman, "Structure-function relations of parasol cells in the normal and glaucomatous primate retina," *Investigative Ophthalmology and Visual Science*, vol. 46, no. 9, pp. 3197–3207, 2005.
- [51] E. Kaplan, B. B. Lee, and R. M. Shapley, "New views of primate retinal function," *Progress in Retinal Research*, vol. 9, pp. 273–336, 1990.
- [52] E. Kaplan and R. M. Shapley, "The primate retina contains two types of ganglion cells, with high and low contrast sensitivity," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 83, no. 8, pp. 2755–2757, 1986.
- [53] E. Kaplan and E. Benardete, "The dynamics of primate retinal ganglion cells," *Progress in Brain Research*, vol. 134, pp. 17–34, 2001.
- [54] B. C. Chauhan and C. A. Johnson, "Test-retest variability of frequency-doubling perimetry and conventional perimetry in glaucoma patients and normal subjects," *Investigative Ophthalmology and Visual Science*, vol. 40, no. 3, pp. 648–656, 1999.
- [55] V. Porciatti, B. Bosse, P. K. Parekh, O. A. Shif, W. J. Feuer, and L. M. Ventura, "Adaptation of the steady-state PERG in early glaucoma," *Journal of Glaucoma*, vol. 23, no. 8, pp. 494–500, 2014.